



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2006

GABA(A) receptor diversity and pharmacology

Möhler, H

Abstract: Because of its control of spike-timing and oscillatory network activity, γ -aminobutyric acid (GABA)-ergic inhibition is a key element in the central regulation of somatic and mental functions. The recognition of GABAA receptor diversity has provided molecular tags for the analysis of distinct neuronal networks in the control of specific pharmacological and physiological brain functions. Neurons expressing 1GABAA receptors have been found to mediate sedation, whereas those expressing 2GABAA receptors mediate anxiolysis. Furthermore, associative temporal and spatial memory can be regulated by modulating the activity of hippocampal pyramidal cells via extrasynaptic 5GABAA receptors. In addition, neurons expressing 3GABAA receptors are instrumental in the processing of sensory motor information related to a schizophrenia endophenotype. Finally, during the postnatal development of the brain, the maturation of GABAergic interneurons seems to provide the trigger for the experience-dependent plasticity of neurons in the visual cortex, with 1GABAA receptors setting the time of onset of a critical period of plasticity. Thus, particular neuronal networks defined by respective GABAA receptor subtypes can now be linked to the regulation of various clearly defined behavioural patterns. These achievements are of obvious relevance for the pharmacotherapy of certain brain disorders, in particular sleep dysfunctions, anxiety disorders, schizophrenia and diseases associated with memory deficits

DOI: <https://doi.org/10.1007/s00441-006-0284-3>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-155970>

Journal Article

Published Version

Originally published at:

Möhler, H (2006). GABA(A) receptor diversity and pharmacology. *Cell and Tissue Research*, 326(2):505-516.

DOI: <https://doi.org/10.1007/s00441-006-0284-3>

GABA_A receptor diversity and pharmacology

H. Möhler

Received: 19 May 2006 / Accepted: 13 June 2006 / Published online: 26 August 2006
© Springer-Verlag 2006

Abstract Because of its control of spike-timing and oscillatory network activity, γ -aminobutyric acid (GABA)-ergic inhibition is a key element in the central regulation of somatic and mental functions. The recognition of GABA_A receptor diversity has provided molecular tags for the analysis of distinct neuronal networks in the control of specific pharmacological and physiological brain functions. Neurons expressing α_1 GABA_A receptors have been found to mediate sedation, whereas those expressing α_2 GABA_A receptors mediate anxiolysis. Furthermore, associative temporal and spatial memory can be regulated by modulating the activity of hippocampal pyramidal cells via extrasynaptic α_5 GABA_A receptors. In addition, neurons expressing α_3 GABA_A receptors are instrumental in the processing of sensory motor information related to a schizophrenia endophenotype. Finally, during the postnatal development of the brain, the maturation of GABAergic interneurons seems to provide the trigger for the experience-dependent plasticity of neurons in the visual cortex, with α_1 GABA_A receptors setting the time of onset of a critical period of plasticity. Thus, particular neuronal networks defined by respective GABA_A receptor subtypes can now be linked to the regulation of various clearly defined behavioural patterns. These achievements are of obvious relevance for the pharmacotherapy of certain brain disorders, in particular sleep dysfunctions, anxiety disorders, schizophrenia and diseases associated with memory deficits.

Keywords Gamma-aminobutyric acid · Benzodiazepines · Anxiety · Learning and memory · Critical period plasticity

Inhibitory interneurons

The activity of inhibitory interneurons, most of which are γ -aminobutyric acid (GABA)-ergic, is thought to set the spatio-temporal conditions required for the various patterns of network oscillations that may be critical for information processing (O'Keefe and Nadel 1978; O'Keefe and Recce 1993; Skaggs et al. 1996; Paulsen and Moser 1998; Engel et al. 2001; Harris et al. 2002; Mehta et al. 2002; Traub et al. 2002; Klausberger et al. 2003).

Diversity of interneurons


To achieve a strict time control of principal cells, GABAergic interneurons display several remarkable features. (1) Their action potential is traditionally faster than that of pyramidal cells and the kinetics of synaptic events that excite inhibitory cells are faster than those that excite pyramidal cells (Martina et al. 1998; Geiger et al. 1997). (2) GABAergic interneurons are morphologically highly diverse, which reflects their multiple functions in neuronal networks (Gupta et al. 2000; Markram et al. 2004; Monyer and Markram 2004). (3) These interneurons show a domain-specific innervation of principal cells; thus, depending on the type of interneuron, particular input domains of pyramidal cells can be selectively regulated and, similarly, the output of pyramidal cells can be specifically regulated by axo-axonic GABAergic interneurons. (4) The response properties of interneuron signalling is shaped by the type of GABA_A receptor expressed synaptically or extrasynaptically. For instance,

H. Möhler (✉)
Institute of Pharmacology and Department of
Chemistry and Applied Biosciences,
University and ETH Zurich,
Winterthurerstrasse 190,
Zürich CH-8057, Switzerland
e-mail: mohler@pharma.unizh.ch

the soma of hippocampal pyramidal cells is innervated by two types of basket cells: the fast spiking parvalbumin-containing basket cells form synapses containing α_1 GABA_A receptors, which display fast kinetics of deactivation (Nyíri et al. 2001; Klausberger et al. 2002; Freund and Buzsáki 1996; Pawelzik et al. 2002), whereas the synapses of the regular spiking cholecystokinin (CCK)-positive basket cells contain α_2 GABA_A receptors, which display slower kinetics than α_1 receptors (Nyíri et al. 2001; Brussaard and Herbison 2000; Hutcheon et al. 2000; Jüttner et al. 2001; Vicini et al. 2001). Axon initial segments of principal cells also contain α_2 receptors, which appear to be kinetically sufficient for simple on/off signalling. Furthermore, distinct GABA_A receptors are segregated to synaptic and extrasynaptic membranes (Nusser et al. 1998; Fritschy and Brünig 2003). Thus, functionally specialized interneurons operate with the kinetically appropriate GABA_A receptor subtypes to regulate network behaviour (Fig. 1, Table 1). Since GABAergic interneurons are operative throughout the brain, a highly diverse repertoire of GABA_A receptors is required.

Retrograde regulation of GABAergic interneurons

Retrograde signalling adds another level of complexity to the regulation of interneuron activity. The terminals of CCK-positive GABAergic basket cells in hippocampus and amygdala contain CB1-cannabinoid receptors (Katona et al. 1999, 2001). These receptors mediate depolarisation-induced suppression of inhibition (DSI) (Pitler and Alger 1994; Alger and Pitler 1995). This phenomenon is due to endocannabinoids which emanate from the postsynaptic cell and act as retrograde signal (Wilson and Nicoll 2001; Maejima et al. 2001). The depolarization of hippocampal pyramidal cells (Pitler and Alger, 1992) and of cerebellar Purkinje cells (Llano et al. 1991) results in a transient decrease in the release of GABA from inhibitory terminals that contain CB1-receptors and synapse onto the depolarized cells (Vincent and Marty, 1993; Pitler and Alger, 1994; Alger and Pitler, 1995). Both the DSI in the hippocampus (Wilson and Nicoll 2001; Maejima et al. 2001) and the cerebellum (Kreitzer and Regehr 2001a,b) are the result of activity-dependent de-novo synthesis and release of endo-



Subunit repertoire

α	1-6
β	1-3
γ	1-3
δ	1
ε	1
θ	1
ρ	1-3

Fig. 1 Representation of a GABAergic synapse depicting major elements of signal transduction. GABA_A receptors are heteromeric membrane proteins linked by an as yet unknown mechanism to the synaptic anchoring protein, gephyrin, and the cytoskeleton. The sequence of subunits corresponds to a modelling proposal (Ernst et al. 2003). The binding sites for GABA and benzodiazepines are located at the interface of the α/β and α/γ_2 subunits, respectively.

Synaptic GABA_A receptors mediate phasic inhibition providing a rapid point-to-point communication for synaptic integration and control of rhythmic network activities. Extrasynaptic GABA_A receptors (not shown) are activated by synaptic spillover or the non-vesicular release of GABA. By mediating tonic inhibition, they provide a maintenance level of reduction in neuronal excitability (Mody and Pearce 2004; Möhler et al. 2005)

Table 1 GABA_A-receptor subtypes (compiled from Möhler et al. 2002; Fritschy and Brünig 2003)

Composition	Pharmacological characteristics	Regional and neuronal localisation	Subcellular localization
$\alpha_1\beta_2\gamma_2$	Major subtype (60 % of all GABA _A receptors). Mediates the sedative, amnestic and, to a large extent, the anticonvulsant action of benzodiazepine site agonists. High affinity for classical benzodiazepines, zolpidem and the antagonist flumazenil	Cerebral cortex (layer 1–VI, selected interneurons and principal cells); hippocampus (selected interneurons and principal cells); pallidum striatum (interneurons); thalamic relay nuclei; olfactory bulb (mitral cells and interneurons); cerebellum (Purkinje cells and granule cells); deep cerebellar nuclei; amygdala; basal forebrain; substantia nigra pars reticulata; inferior colliculus; brainstem	Synaptic (soma and dendrites) and extrasynaptic in all neurons with high expression
$\alpha_2\beta_3\gamma_2$	Minor subtype (15–20%). Mediates anxiolytic action of benzodiazepine site agonists. High affinity for classical benzodiazepine agonists and the antagonist flumazenil. Intermediate affinity for zolpidem	Cerebral cortex (layers I–IV) hippocampal formation (principal cells mainly on the axon initial segment); olfactory bulb (granule cells); striatum (spiny stellate cells); inferior olivary neurons (mainly on dendrites); hypothalamus; amygdala (principal cells); superior colliculus; motor neurons	Mainly synaptic, enriched in axon initial segment of cortical and hippocampal pyramidal cells
$\alpha_3\beta_n\gamma_2$	Minor subtype (10–15%). High affinity for classical benzo-diazepine agonists and the antagonist flumazenil. Intermediate affinity for zolpidem	Cerebral cortex (principal cells in particular in layers V and VI; some axon initial segments); hippocampus (some hilar cells); olfactory bulb (tufted cells); thalamic reticular neurons; cerebellum (Golgi type II cells); medullary reticular formation; inferior olivary neurons; amygdala; superior colliculus; brainstem; spinal cord; medial septum; basal forebrain cholinergic neurons; raphe and locus coeruleus (serotonergic and catecholaminergic neurons)	Mainly synaptic, including some axon initial segments; extrasynaptic in inferior olivary neurons
$\alpha_4\beta_n\delta$	Less than 5 % of all receptors. Insensitive to classical benzodiazepine agonists and zolpidem	Dentate gyrus (granule cells); thalamus	Extrasynaptic (no direct morphological evidence)
$\alpha_5\beta_3\gamma_2$	Less than 5% of all receptors; high affinity for classical benzodiazepine agonists and the antagonist flumazenil. Low affinity for zolpidem	Hippocampus (pyramidal cells); olfactory bulb (granule cells, periplomerular cells); cerebral cortex; amygdala; hypothalamus; superior colliculus; superior olivary neurons; spinal trigeminal neurons; spinal cord	Extrasynaptic in hippocampus, cerebral cortex, and olfactory bulb; synaptic and extrasynaptic in spinal trigeminal nucleus and superior olivary nucleus
$\alpha_6\beta_{2,3}\gamma_2$	Less than 5 % of all receptors. Insensitive to classical $\alpha_6\beta_{2,3}\delta$ benzodiazepine agonists and zolpidem. Minor population. Lacks benzodiazepine site	Cerebellum (granule cells); dorsal cochlear nucleus	Synaptic (cerebellar glomeruli) and extrasynaptic on granule cell dendrites and soma

cannabinoids from the postsynaptic neuron. By interacting with CB1 receptors, the calcium influx into the presynaptic terminal is reduced (Kreitzer and Regehr 2001a; Caulfield and Brown 1992) resulting in a decrease of GABA release. Pharmacologically, the inhibition of the degradation of anandamide resulted in an CB1 receptor-mediated anxiolytic response (Kathuria et al. 2003). Recently, endocanna-

binoids were shown to mediate not only transient, but also long term changes in inhibitory synaptic transmission. Endocannabinoid production, stimulated through metabotropic glutamate receptor activation in hippocampal pyramidal cells, caused a long lasting reduction in GABAergic signaling onto the pyramidal cells (Chevalleyre and Castillo 2003). A similar change in long-term synaptic plasticity of

the GABAergic system was observed in the amygdala. The release of endocannabinoids in the basolateral amygdala contributed to the extinction of aversive memory based on a long lasting decrease of GABAergic signalling (Marsicano et al. 2002). Thus, CB1 receptor activation in the hippocampus, amygdala and possibly other parts of the brain result in reduced levels of anxiety. This is due to either a transient depression of GABA release or a modulation of long term plasticity at the respective synapses. In addition, endocannabinoids act as retrograde signals at excitatory glutamatergic synapses where they mediate a depolarisation-induced suppression of excitation (Kreitzer and Regehr 2001a) which may also contribute to their behavioral effects.

Structure of GABA_A receptors

Like other members of the nicotinic superfamily of ligand-gated ion channels, ionotropic GABA receptors are considered to consist of 5 protein subunits arranged around a central pore that constitutes the actual ion channel. Each subunit has a large extracellular N-terminal domain which incorporates part of the agonist/antagonist binding site, followed by three membrane spanning domains (M1–3), an intracellular loop of variable length and a fourth membrane spanning domain (M4), with the C-terminal end being extracellular. Each subunit arranges itself such that the second membrane-spanning domain (M2) forms the wall of the channel pore. The cytoplasmic loop, between the third and fourth transmembrane domains (M3 and M4), is believed to be the target for protein kinases, required for subcellular targeting and membrane clustering of the receptor. There are 16 different subunits comprising the GABA_A receptor family: α_{1-6} , β_{1-3} , γ_{1-3} , δ , ϵ , π and θ . In addition, there are splice variants of many of these subunits. Based on this subunit repertoire more than 2000 different GABA_A receptors could exist if the subunit combinations were restricted to those containing two α , two β and one other subunit. In fact, studies of native GABA_A receptors suggest that there may be less than 20 widely occurring GABA_A receptor subtype combinations, with the major combinations being $\alpha_1\beta_2\gamma_2$, $\alpha_2\beta_3\gamma_2$, $\alpha_3\beta_3\gamma_2$ (for review Barnard et al. 1998; Whiting et al. 2000; Sieghart and Sperk 2002; Möhler et al. 2000, 2002, 2005; Möhler 2001, 2002; Fritschy et al. 2004; Rudolph and Möhler 2006) (Table 1).

In the retina homomeric receptors consisting of the ρ subunit represent a particular class of GABA gated chloride channels. Their GABA site is insensitive to bicuculline and baclofen and they are not modulated by barbiturates or benzodiazepines. Due to these distinctive features the receptors are sometimes termed GABA_c-receptors

(Bormann 2000), although they are a homomeric class of GABA_A-receptors (Barnard et al. 1998).

The physiological significance of the structural diversity of GABA_A receptors lies in the provision functionally diverse of receptors which differ in their channel kinetics, affinity for GABA, rate of desensitization and ability for transient chemical modification such as phosphorylation. In addition, GABA_A receptor subtypes differ in their topology in neuronal networks in that they show a cell-type specific expression and-in case of multiple receptor subtypes present in a neuron-a domain-specific location (Table 1).

Pharmacology of GABA_A receptors

Diazepam-sensitive GABA_A receptors

Receptors containing the α_1 , α_2 , α_3 or α_5 subunits in combination with any of the β subunits and the γ_2 subunit are most prevalent in the brain. These receptors are sensitive to benzodiazepine modulation. The major receptor subtype is assembled from the subunits $\alpha_1\beta_2\gamma_2$, with only a few brain regions lacking this receptor (e.g. granule cell layer of the olfactory bulb, reticular nucleus of the thalamus, spinal cord motoneurons) (Fritschy and Mohler 1995; Pirker et al. 2000, Fritschy and Brünig 2003).

Receptors containing the α_2 or α_3 subunit are considerably less abundant and are highly expressed in brain areas where the α_1 subunit is absent or present at low levels. The α_2 and α_3 subunits are frequently coexpressed with the β_3 and γ_2 subunits which is particularly evident in hippocampal pyramidal neurons ($\alpha_2\beta_3\gamma_2$) and in cholinergic neurons of the basal forebrain ($\alpha_3\beta_3\gamma_2$). The α_3 GABA_A receptors are the main subtypes expressed in monoaminergic and basal forebrain cholinergic cells (Gao et al. 1993) and are, in addition, strategically located in the thalamic reticular nucleus for modulating the thalamo-cortical circuit (Huntsmann et al. 1999). Marked differences in desensitization kinetics have been reported between synaptic α_2 - and extrasynaptic α_3 -receptors whereby the latter desensitize very slowly (Devor et al. 2001). The factors regulating GABA_A receptor kinetics at synaptic and extrasynaptic sites are yet unknown (Moss and Smart 2001). The ligand-binding profile of the α_2 - and α_3 -receptors differs from that of $\alpha_1\beta_2\gamma_2$ by having a considerably lower displacing potency for ligands such as β CCM, CL 218,872, and zolpidem.

Receptors containing the α_5 subunit are of minor abundance in the brain but are expressed to a significant extent in the hippocampus, where they comprise 15 % to 20 % of the diazepam-sensitive GABA_A receptor population, predominately coassembled with the β_3 and γ_2 subunits. Pharmacologically, the α_5 -receptors are differentiated from

$\alpha_1\beta_2\gamma_2$, $\alpha_2\beta_3\gamma_2$ and $\alpha_3\beta_3\gamma_2$ receptors by a lower affinity to CL 218, 872 and near-insensitivity to zolpidem.

The subunits γ_1 and γ_3 characterize a small population of receptors that contain various types of α and β subunits. Due to their reduced affinity for the classical benzodiazepines they do not appear to contribute to any great extent to their pharmacology in vivo.

It should be kept in mind that complex benzodiazepine actions such as the development of tolerance can implicate more than a single receptor subtype. For instance, while the sedative action of diazepam is mediated by α_1 GABA_A receptors (see below), the development of tolerance to this action under chronic diazepam treatment requires the interaction with both α_1 GABA_A receptors and α_5 GABA_A receptors (van Rijnsoever et al. 2004).

Diazepam-insensitive GABA_A receptors

GABA_A receptors that do not respond to clinically used ligands such as diazepam, flunitrazepam, clonazepam, and zolpidem are of low abundance in the brain and are largely characterized by the α_4 and α_6 subunits. Receptors containing the α_4 subunit are generally expressed at very low abundance but more prominently in thalamus and dentate gyrus (Pirker et al. 2000); those containing the α_6 subunit are restricted to the granule cell layer of the cerebellum (about 30–50 % of all GABA_A receptors in the cerebellum; Nusser et al. 1996; Pölzl et al. 2003). Both receptor populations are structurally heterogeneous, and the majority of the α_6 -containing receptors contain the γ_2 subunit in the $\alpha_6\alpha_1\beta_2\gamma_2$ subunit combination (Pölzl et al. 2003). Apart from the lack of affinity of classical benzodiazepines, the benzodiazepine-site profile of α_4 and α_6 receptors is characterized by a low affinity for flumazenil and bretazenil and an agonistic efficacy of Ro 15-4513 and bretazenil (Benson et al. 1998). The δ subunit is frequently coassembled with the α_4 or the α_6 subunit in benzodiazepine insensitive receptors (Möhler et al. 2000; Whiting et al. 2000; Möhler 2001). Receptors containing the δ subunit are located exclusively at extrasynaptic sites as shown in dentate gyrus and cerebellum. They are tailor made for tonic inhibition, due to their high affinity for GABA and slow desensitization kinetics (Brickley et al. 1996, 2001).

GABA_A receptor subtypes: a new pharmacology

The selective pattern of expression of GABA_A receptor subtypes opened the possibility to modulate distinct neuronal circuits, provided novel ligands were found which displayed a differential interaction with GABA_A receptor subtypes based on either selective affinity or selective efficacy (Table 2). Such agents would be expected to display therapeutic indications which are more selective

than those of the classical benzodiazepines and go beyond their spectrum of activity.

The dissection of receptor pharmacology has been achieved experimentally by generating four lines of point-mutated mice in which the receptors containing the α_1 , α_2 , α_3 or α_5 subunits, respectively, are rendered diazepam-insensitive by replacing a conserved histidine residue (H) in the drug-binding domain by an arginine (R; Benson et al. 1998; Rudolph et al. 1999; Löw et al. 2000; Crestani et al. 2002). In the respective point-mutated mice, the pharmacological action linked to the point-mutated receptor should be missing and thereby reveal the pharmacological relevance of the respective receptor in wild-type mice. Since the subunit composition and distribution of GABA_A receptor subtypes are largely conserved between rodents and non-human primates, the results are thought to be relevant to the human condition (Rudolph and Möhler 2004).

GABA_A receptors for sedation and sleep

Frequently, sedation is taken as a surrogate marker for hypnotic activity. The sedative component of benzodiazepines, measured by the reduction of locomotor activity, has been attributed to neuronal circuits expressing α_1 GABA_A receptors, the most prevalent receptor subtype in the brain. Mice in which the α_1 GABA_A receptor has been rendered diazepam-insensitive by a point mutation [α_1 (H101R)] fail to be sedated by diazepam (Rudolph et al. 1999; McKernan et al. 2000). Ligands with a preferential affinity for α_1 receptors such as zolpidem or zaleplon are used as hypnotics (Table 2). Similarly, the changes in the electroencephalogram (EEG) pattern induced by zolpidem in wild-type mice are almost exclusively mediated via α_1 GABA_A receptors (Kopp et al. 2004a). However, the changes in sleep architecture (suppression of rapid eye movement or REM sleep) and EEG-frequency profiles (reduction of slow-wave sleep, increase in fast β -frequencies) induced by classical benzodiazepines are largely attributable to effects mediated by receptors others than α_1 (Tobler et al. 2001). The enhancement of α_2 GABA_A receptors by diazepam appears to have the most pronounced effect on the sleep EEG in wild-type mice. When the α_2 GABA_A receptor is rendered diazepam-insensitive by a point mutation [α_2 (H101R)], the diazepam-induced suppression of δ -waves, the increase in fast β -waves in non-REM sleep (>16 Hz) and the diazepam-induced increase of θ -waves in REM sleep are strongly attenuated (Kopp et al. 2004b). Thus, the hypnotic EEG fingerprint of diazepam can be dissociated from its sedative action. Future hypnotics might target changes in the EEG pattern, which are characteristic of physiological sleep, and thereby aim at improving sleep quality. For instance, the GABA-mimetic gaboxadol (4,5,6,7-tetrahydroisoxazolo[5,4-c]pyridine-3-ol hydrochloride)

Table 2 GABA_A receptor subtype ligands*

Drug	Main activity	Interaction with recombinant GABA _A receptors ^{1,2}	Reference
A. Benzodiazepine site ligands			
Zolpidem	Hypnotic	Preferential affinity for α_1	Dämgen and Lüddens, 1999
Zaleplone	Hypnotic	Preferential affinity for α_1	Dämgen and Lüddens, 1999
Indiplon	Hypnotic	Preferential affinity for α_1	Foster et al. 2004
L-838 417	Anxiolytic	Comparable affinity at α_1 , α_2 , α_3 , α_5 subtype. Partial agonist at α_2 , α_3 , α_5 (not α_1) subtype	McKernan et al. 2000
Ocinaplon	Anxiolytic	Comparable affinity at α_1 , α_2 , α_3 , α_5 subtype. Partial agonist at α_2 , α_3 , α_5 subtype nearly full agonist at α_1	Lippa et al. 2005
SL 651 498	Anxiolytic	Agonist at α_2 , α_3 , partial agonist at α_1 and α_5 subtype	Griebel et al. 2003
TPA 023	Anxiolytic	Partial agonist at α_2 , α_3 subtypes, antagonist at α_1 , α_5 subtypes	Atack et al. 2006a
TPA 003	Anxiolytic	Partial agonist at α_3 subtype	Dias et al. 2005
ELB 139	Anxiolytic	Selective receptor profile uncertain	Langen et al. 2005
L-655 708	Memory enhancer, Anxiogenic	Partial inverse agonist with preference for α_5 subtype	Sternfeld et al. 2004; Chambers et al. 2004; Navarro et al. 2002, 2004
α_3 IA	Anxiogenic	Weak inverse agonist at α_3	Atack et al. 2005
B. Modulatory site other than benzodiazepine site			
Ethanol	Anxiolytic Sedative	High sensitivity (≥ 3 mM) at $\alpha_4(\alpha_6)\beta_{3\delta}^3$; Medium sensitivity (≥ 30 mM) at $\alpha_4(\alpha_6)\beta_{2\delta}^3$; Low sensitivity (≥ 100 mM) at $\alpha_4(\alpha_6)\beta_{3\gamma_2}$	Wallner et al. 2003
Neurosteroids (e.g. 3 α , 5 α -THDOC)	Anxiolytic Sedative Anaesthetic	High sensitivity at δ -containing subtypes ³ and at α_1 , α_3 receptors in combination with β_1	Belelli and Lambert 2005
Intravenous anaesthetics (Etomidate Propofol)	Sedative Anaesthetic	Act on receptor subtypes containing β_3 i.e. mainly α_2 and α_3 subtypes	Rudolph and Antkowiak 2004
C. GABA site			
Gaboxadol	Hypnotic	Partial agonist at α_1 , α_3 subtypes full agonist at α_5 and superagonist at $\alpha_4\beta_3\delta$ receptors ²	Stornstovu and Ebert 2003

*This table is a modified version from Rudolph and Möhler (2006)

¹Classical partial agonists that do not differentiate between GABA_A receptor subtypes such as Bretazenil (Haefely et al. 1990) or Pagoclone (Atack et al. 2006b) are not considered in this review.

²Data should be treated with caution as properties of recombinant receptors that are expressed in foreign host cells might not give an accurate reflection of their neuronal counterparts.

³GABA is a weak partial agonist on δ -containing receptors, which largely explains the strong modulatory response of ligands acting on δ -containing receptors (Bianchi and MacDonald 2003). THDOC, 5 α -pregnane-3 α ,21-diol-20-one.

ride; synonym THIP), which interacts preferentially with $\alpha_4\beta_3\delta$ GABA_A receptors in vitro (Brown et al. 2002; Stornstovu and Ebert 2003) has been found to enhance slow-wave sleep in vivo (Lancel and Steiger 1999; Huckle 2004).

GABA_A receptors for anxiolytic action

Since α_1 GABA_A receptors were found to mediate sedation but not anxiolysis (Rudolph et al. 1999; McKernan et al. 2000), the anxiolytic activity of benzodiazepines was

expected to reside in one or several of the remaining benzodiazepine-sensitive GABA_A receptors (α_2 , α_3 , α_5). The differentiation of GABA_A receptors by knock-in point mutations showed that the α_2 receptor, but not the α_3 or α_5 GABA_A receptor, mediated the anxiolytic activity of diazepam (Löw et al. 2000; Crestani et al. 2002). In α_2 (H101R) mice, but not in α_3 (H126R) or α_5 (H105R) mice, diazepam failed to induce anxiolytic activity (light-dark paradigm, elevated plus maze). With respect to the α_2 GABA_A receptor, a highly selective target for the anxiolytic activity of benzodiazepine tranquillizers had

been identified. In keeping with this notion, the benzodiazepine site ligand L-838417, which exhibited efficacy at α_2 , α_3 and α_5 but not α_1 GABA_A receptors, proved to be anxiolytic in wild-type rats (Table 2; McKernan et al. 2000). Similarly, partial agonists of 3-heteroaryl-2-pyridones acting at the benzodiazepine site with efficacy at α_2 , α_3 and α_5 receptors, but not at α_1 receptors, were found to show anxiolytic activity in rodents (Table 2; Collins et al. 2002). Nevertheless, the extent to which the α_3 GABA_A receptor component contributed to the anxiolytic activity of these ligands remained to be clarified. In mice that lacked α_3 GABA_A receptors, the anxiolytic activity of diazepam was undiminished (Yee et al. 2005). However, an α_3 -selective inverse agonist was anxiogenic and proconvulsant in rodents (Table 2; Collins et al. 2002). In addition, TP003 with selective efficacy at the α_3 GABA_A receptor was anxiolytic, although only at high receptor occupancy (Dias et al. 2005). Classical benzodiazepines exert anxiolysis at low receptor occupancy suggesting that the α_2 GABA_A receptors, and not the α_3 GABA_A receptors, are the major mediators of this activity. The contribution of α_3 GABA_A receptors is unlikely to be of major relevance. Thus, the strategy to develop novel daytime anxiolytics, which are free of sedation, is clear (Whiting 2003; Möhler et al. 2002; Möhler et al. 2005).

The α_2 GABA_A receptors by their preponderant localization on the axon-initial segment of principal cells in the cerebral cortex and hippocampus can control the output of these cells. In addition, among α_1 , α_2 and α_3 GABA_A receptors, α_2 receptors are prominent in the central nucleus of the amygdala, a key area for the control of emotions, whereas α_1 GABA_A receptors are totally absent (Marowsky et al. 2004). Thus, by their strategic distribution in brain areas involved in anxiety responses, α_2 GABA_A receptors represent key substrates for anxiolytic drug action.

GABA_A receptors for learning and memory

Hippocampal pyramidal cells express various structurally diverse GABA_A receptors in a domain-specific manner. Whereas α_1 and α_2 GABA_A receptors are largely synaptic, α_5 GABA_A receptors are located extrasynaptically at the base of the spines and on the adjacent shaft of the pyramidal cell dendrite. The α_5 GABA_A receptors are therefore in a privileged position to modulate the excitatory input arising at the spines via N-methyl-D-aspartate (NMDA) receptors. The introduction of a point mutation (H105R) into the α_5 subunit is associated with a specific reduction of the hippocampal α_5 -subunit-containing GABA_A receptors, whereas the pattern of distribution is undisturbed (Crestani et al. 2002). Mice with a partial deficit of α_5 GABA_A receptors in the hippocampus show an improved performance in trace fear conditioning, a hippo-

campus-dependent memory task (Crestani et al. 2002). In addition, these mutants display a resistance to the extinction of conditional fear over several days (Yee et al. 2004). Similarly, in a mouse line in which α_5 GABA_A receptors are absent in the entire brain (Collinson et al. 2002; Whiting 2003), improved performance has been observed in the water maze model of spatial learning. Furthermore, a partial inverse agonist acting at α_5 GABA_A receptors enhances the performance of wild-type rats in the water maze test (Chambers et al. 2003; Table 2). Thus, neuronal inhibition in the hippocampus mediated via α_5 GABA_A receptors is a critical element in the regulation of the acquisition and expression of associative memory. Of note, the behavioural consequences of an impairment of α_5 GABA_A receptors are opposite to those of an NMDA receptor deficit as shown in spatial and temporal associative memory tasks. Thus, these two receptor systems seem to play a complementary role in controlling signal transduction at the hippocampal principal cells.

GABA_A receptors for sensorimotor processing

A deficit in GABAergic inhibitory control is one of the major hypothesis underlying the symptomatology of schizophrenia (Lewis et al. 2005). A potential contribution of GABA_A receptor subtypes has therefore been investigated with regard to the overactivity of the dopaminergic system, an overactivity considered to be a major factor in schizophrenia. The dopaminergic system is under GABAergic inhibitory control mainly via α_3 -containing GABA_A receptors (Fritschy and Möhler 1995; Pirker et al. 2000). Their functional role has been explored in mice lacking the α_3 subunit gene. Mice with an α_3 -knockout display no adaptive changes in the expression of α_1 , α_2 and α_5 subunits and their anxiety-related behaviour is normal. However, they exhibit a marked deficit in prepulse inhibition of the acoustic startle reflex, pointing to a deficit in sensorimotor information processing (Yee et al. 2005). This deficit in prepulse inhibition is normalized by the administration of the antipsychotic dopamine D2-receptor antagonist haloperidol, suggesting that the behavioural phenotype is caused by hyperdopaminergia (Yee et al. 2005). Attenuation of prepulse inhibition is a frequent phenotype of psychiatric conditions, including schizophrenia. These results suggest that α_3 -selective agonists might constitute an effective treatment for sensorimotor gating deficits in various psychiatric conditions, a view supported by the observation that the partial benzodiazepine-site agonist bretazenil, in earlier open clinical trials, displayed an antipsychotic activity similar to neuroleptic drugs (Delini-Stula and Berdah Tordjman 1996). The α_3 -selective agonists might lack the sedative or extrapyramidal side-effects of classical neuroleptics and would thus be valuable agents.

Among various brain structures, the hippocampus is believed to play an important role in the modulation of prepulse inhibition. In α_5 (H105R) point-mutated mice, the expression of the α_5 subunit containing GABA_A receptors in the hippocampus is reduced (see above; Crestani et al. 2002). In these animals, prepulse inhibition is attenuated concomitant with an increase in spontaneous locomotor activity (Hauser et al. 2005). Thus, the α_5 -subunit-containing GABA_A receptors, which are located extrasynaptically and are thought to mediate tonic inhibition, are important regulators of the expression of prepulse inhibition and locomotor exploration. Post-mortem analyses of brains from patients with schizophrenia have consistently revealed structural abnormalities of developmental origin in the hippocampus (Lewis et al. 2005). Such abnormalities may include disturbances of α_5 GABA_A receptor function given that schizophrenic patients are known to exhibit a deficit in prepulse inhibition. Thus, agonists acting on both α_3 and α_5 GABA_A receptors may be beneficial in overcoming this endophenotypic manifestation of the disease.

GABA_A receptors for anaesthetic action

The targets that mediate the clinical effects of general anaesthetics are largely unknown (Campagna et al. 2003; Rudolph and Antkowiak 2004). Recent work has focused on the role of GABA_A receptors, with studies being based on the analysis of point-mutated knock-in mice carrying point mutations in the β_3 and β_2 subunits of the GABA_A receptor. These mutations render the GABA_A receptors containing the respective subunits insensitive to modulation by etomidate, propofol and certain volatile anaesthetics, such as enflurane. The β_3 -containing GABA_A receptors have been found to mediate the immobilizing action of etomidate and propofol apparently in full (Table 2; Jurd et al. 2003) and of enflurane, isoflurane and halothane in part (Jurd et al. 2003; Lambert et al. 2005; Liao et al. 2005). In addition, they mediate part of the hypnotic action of etomidate and propofol (Jurd et al. 2003) but apparently not of the volatile anaesthetics (Jurd et al. 2003; Lambert et al. 2005). In contrast, the hypnotic action of etomidate is mediated by β_2 -containing GABA_A receptors (Reynolds et al. 2003). Further studies have revealed that the respiratory depressant action of etomidate and propofol is also mediated by β_3 -containing GABA_A receptors, whereas the heart-rate depressant action and, to a large part, the hypothermic action of etomidate and propofol are mediated by other targets (Zeller et al. 2005; Cirone et al. 2004). Thus, a β_3 -selective agent would be predicted to be immobilizing and respiratory depressant but would largely lack the heart-rate depressant and hypothermic actions of etomidate and propofol. The analysis of α subunits involved in mediating the actions of general anaesthetics

is expected to result in further insights into the contribution of GABA_A receptors to anaesthesia. Mutations in α subunits have been identified in recombinant studies that render $\alpha_x\beta_x\gamma_2$ GABA_A receptors insensitive to specific volatile anaesthetics but not to etomidate or propofol (Mihic et al. 1997; Krasowski et al. 1998). Studies with knock-in mice carrying these mutations are expected to yield information as to the contribution of individual GABA_A receptors subtypes and the GABA_A receptor family as a whole to the action of volatile general anaesthetics.

GABA_A receptors controlling postnatal development

Apart from its trophic role in embryonal development (Represa and Ben-Ari 2005), GABA is a major determinant for postnatal developmental plasticity; this has been investigated in various sensory systems. For instance, in the rodent somatosensory system, axons from each whisker form a somatotopic map in the cortex, known as the barrel map. During a critical period of neonatal development, this barrel map is fine-tuned in response to sensory experience based on a variety of synaptic mechanisms involving not only excitatory, but also inhibitory circuits (Foeller and Feldmann 2004). The role of inhibitory circuits in synaptic reorganization is similarly apparent in the auditory system. Recent work has revealed a dramatic remodelling of inhibitory synapses shortly after the onset of hearing (aural dominance bands). The restructuring relies on both spontaneous and sensory-evoked neural activity (Kandler 2004). In the postnatal visual system, the role of GABA has been investigated in detail.

In the visual system, developmental plasticity is most apparent in the formation of ocular dominance columns in layer IV of the primary visual cortex. Cortical territories receiving neuronal input from one eye alternate with territories from the other eye. Initially, at birth, the thalamic inputs from both eyes to the visual cortex are totally overlapping (Ferster 2004). The separation of the visual inputs into ocular dominance columns only arises in the subsequent phase of remodeling. This process is sensitive to light as shown by classical work on the influence of monocular deprivation on ocular dominance plasticity (Wiesel and Hubel 1963). After the closure of one eye during a critical period of early postnatal life, the input from the open eye subsequently has a larger cortical territory than the input from the deprived eye. Critical period plasticity is best viewed as a continuum of local circuit computations that result in a structural rewiring of thalamic afferents (Hensch 2005).

The mechanism of visual cortical plasticity was analyzed in detail with regard to the contribution of intracortical

GABAergic transmission. GABAergic transmission was modulated locally by infusion of the benzodiazepine agonist diazepam or the inverse agonist DMCM. Following chronic infusion of diazepam into the striate cortex (starting at postnatal days 14–17), the spacing of the ocular columns was widened, whereas infusion of DMCM reduced the spacing (Hensch and Stryker 2004). Visual responsiveness remained undisturbed under these conditions (Hensch and Stryker 2004). Thus, intracortical GABA interneurons shape the geometry of the incoming thalamic arbours. In addition, the degree of GABAergic inhibition has been found to be a key determinant for the onset of critical period plasticity. The enhancement of GABA transmission by diazepam has long been known to induce the premature onset of the critical period (Fagiolini and Hensch 2000). It has now been found that only circuits containing α_1 GABA_A receptors drive cortical plasticity, whereas α_2 -enriched connections separately regulate neuronal firing (Fagiolini et al. 2004). These results are based on the use of knock-in mice in which the respective individual α -subunit had been rendered diazepam-insensitive by a point mutation (Rudolph et al. 1999; Löw et al. 2000). These recent findings therefore present a cellular and molecular basis for critical period plasticity in the visual cortex triggered by neuronal inhibition (Fagiolini et al. 2004; Ferster 2004).

For ocular stripes to form postnatally, activity in nearby inputs from the same eye are considered to cooperate with each other as cluster of cortical cells in their bid to take over cortical territory (Ferster 2004). By lateral GABAergic inhibitory connections, activity in more distant cells must be “anti-correlated”. Inputs from the same eye are therefore suppressed in their bid to take over the adjacent territories. The pattern of ocular dominance columns thus arises during the segregation of eye-specific inputs to the visual cortex in a self-organizing process. The cortex itself, through a specific type of GABAergic interneuron, plays a central role in organizing this pattern (Ferster 2004). The special developmental function of neocortical α_1 GABA_A receptors suggests that constraints should be placed on drugs designated for use in human infants (Fagiolini et al. 2004).

References

- Alger BE, Pitler TA (1995) Retrograde signaling at GABA_A-receptor synapses in the mammalian CNS. *Trends Neurosci* 18:333–340
- Atack JR, Hutson PH, Collinson N, Marchall G, Bentley G, Moyes C, Cook SM, Collins I, Wafford K, McKernan RM, Dawson GR (2005) Anxiogenic properties of an inverse agonist selective for α_3 subunit-containing GABA_A receptors. *Br J Pharmacol* 144:357–366
- Atack JR, Wafford K, Tye SJ, Cook S, Sohal B, Pike A, Sur C, Melillo D, Bristow L, Bromidge F, et al (2006a) TPA023 an agonist selective for α_2 - and α_3 -containing GABA_A receptors, is a non-sedating anxiolytic in rodents and primates. *J Pharm Exp Ther* 316:410–422
- Atack JR, Wafford K, Tye SJ, Cook S, Sohal B, Pike A, Sur C, Melillo D, Bristow L, Bromidge F, et al (2006b) The in vivo properties of pagoclone in rat are most likely mediated by 5'-hydroxy pagoclone. *Neuropharmacology* (in press)
- Barnard EA, Skolnick P, Olsen RW, Möhler H, Sieghart W, Biggio G, Braestrup C, Bateson AN, Langer SZ (1998) Subtypes of γ -aminobutyric acid_A receptors: classification on the bases of subunit structure and receptor function. *Pharmacol Rev* 50:291–313
- Belelli D, Lambert JJ (2005) Neurosteroids: endogenous regulators of the GABA(A) receptor. *Nat Rev Neurosci* 6:565–575
- Benson J, Löw K, Keist R, Möhler H, Rudolph U (1998) Pharmacology of recombinant GABA_A receptors rendered diazepam-insensitive by point-mutated α -subunits. *FEBS Lett* 431:400–404
- Bianchi MT, McDonald RL (2003) Neurosteroids shift partial agonist activation of GABA(A) receptor channels from low- to high-efficacy gating patterns. *J Neurosci* 23:10934–10943
- Bormann J (2000) The “ABC” of GABA receptors. *Trends Pharmacol Sci* 21:16–19
- Brickley SG, Cull-Candy SG, Farrant M (1996) Development of a tonic form of synaptic inhibition in rat cerebellar granule cells resulting from persistent activation of GABA_A receptors. *J Physiol (Lond)* 497:753–759
- Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M (2001) Adaptive regulation of neuronal excitability by a voltage independent potassium conductance. *Nature* 409:88–92
- Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA (2002) Pharmacological characterization of a novel cell line expressing human $\alpha 4\beta 3\delta$ GABA_A receptors. *Br J Pharmacol* 136:965–974
- Brussaard AB, Herbison AE (2000) Long-term plasticity of postsynaptic GABA_A-receptor function in the adult brain: insights from the oxytocin neurone. *Trends Neurosci* 23:190–195
- Campagna JA, Miller KW, Forman SA (2003) Mechanisms of actions of inhaled anesthetics. *N Engl J Med* 348:2110–2124
- Caulfield MP, Brown DA (1992) Cannabinoid receptor agonists inhibit Ca current in NG108-15 neuroblastoma cells via a pertussis toxin-sensitive mechanism. *Br J Pharmacol* 106:231–232
- Chambers MS, Attack JR, Broughton HB, Collinson N, Cook S, Dawson GR, Hobbs SC, Marshall G, Maubach KA, Pillai GV, Reeve AJ, MacLeod AM (2003) Identification of a novel, selective GABA_A α_5 receptor inverse agonist which enhances cognition. *J Med Chem* 46:2227–2240
- Chambers MS, Attack JR, Carling RW, Collinson N, Cook SM, Dawson GR, Ferris P, Hobbs SC, O'Connor D, Marshall G et al (2004) An orally bioavailable, functionally selective inverse agonist at the benzodiazepine site of GABA_A α_5 receptors with cognition enhancing properties. *J Med Chem* 47:5829–5832
- Chevalleyre V, Castillo PE (2003) Heterosynaptic LTD of hippocampal GABAergic synapses: a novel role of endocannabinoids in regulating excitability. *Neuron* 38:461–472
- Cirone J, Rosahl TW, Reynolds DS, Newman RJ, O'Meara GF, Hutson PH, Wafford KA (2004) Gamma-aminobutyric acid type A receptor beta 2 subunit mediates the hypothermic effect of etomidate in mice. *Anesthesiology* 100:1438–1445
- Collins I, Moyes C, Davey WB, Rowley M, Bromidge FA, Quirk K, et al (2002) 3-Heteroaryl-2-pyridones: benzodiazepine site ligands with functional selectivity for α_2/α_3 -subtypes of human GABA(A) receptor-ion channels. *J Med Chem* 45:1887–1900
- Collinson N, Kuenzi FM, Jarolimek W, Maubach KA, Cothliff R, Sur C, et al (2002) Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the α_5 subunit of the GABA_A receptor. *J Neurosci* 22:5572–5580

- Crestani F, Keist R, Fritschy JM, Benke D, Vogt K, Prut L, Bluethmann H, Möhler H, Rudolph U (2002) Trace fear conditioning involves hippocampal α_5 GABA_A receptors. *Proc Natl Acad Sci USA* 99:8980–8985
- Dämgen K, Lüddens H (1999) Zaleplon displays a selectivity to recombinant GABA_A receptors different from zolpidem, zopiclone and benzodiazepines. *Neurosci Res Comm* 25: 139–148
- Dellini-Stula A, Berdah-Tordjman D (1996) Antipsychotic effects of bretazenil, a partial benzodiazepine agonist in acute schizophrenia—a study group report. *J Psychiatr Res* 30:239–250
- Devor A, Fritschy JM, Yarom Y (2001) Spatial distribution and subunit composition of GABA_A receptors in the inferior olivary nucleus. *J Neurophysiol* 85:1686–1696
- Dias R, et al (2005) Evidence for a significant role of α_3 -containing GABA_A receptors in mediating the anxiolytic effects of benzodiazepines. *J Neurosci* 25:10682–10688
- Engel AK, Fries P, Singer W (2001) Dynamic predictions: oscillations and synchrony in top-down processing. *Nat Rev Neurosci* 2:704–716
- Ernst M, Brauchart D, Boresch S, Sieghart W (2003) Comparative modeling of GABA_A receptors: limits, insights, future developments. *J Neuroscience* 4:933–943
- Fagioli M, Hensch T (2000) Inhibitory threshold for critical-period activation in primary visual cortex. *Nature* 404:183–186
- Fagioli M, Fritschy JM, Löw K, Möhler H, Rudolph U, Hensch T (2004) Specific GABA_A circuits for visual cortical plasticity. *Science* 303:1681–1683
- Ferster D (2004) Blocking plasticity in the visual cortex. *Science* 303:1619–1621
- Foeller E, Feldmann DE (2004) Synaptic basis for developmental plasticity in somatosensory cortex. *Curr Opin Neurobiol* 14: 89–95
- Foster AC, Pellemounter MA, Cullen MJ, Lewis D, Joppa M, Chen TK, Bozigian HP, Gross RS, Gogas KR (2004) In vivo pharmacological characterization of indiplon, a novel pyrazolopyrimidine sedative-hypnotic. *J Pharmacol Exp Ther* 311: 547–559
- Freund TF, Buzsaki G (1996) Interneurons of the hippocampus. *Hippocampus* 6:345–470
- Fritschy JM, Brünig I (2003) Formation and plasticity of GABAergic synapses: physiological mechanisms and pathophysiological implications. *Pharmacol Ther* 98:299–323
- Fritschy JM, Möhler H (1995) GABA_A receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol* 359:154–194
- Fritschy JM, Crestani F, Rudolph U, Möhler H (2004) GABA_A receptor subtypes with special reference to memory function and neurological disorders. In: Hensch TK, Fagioli M (eds) *Excitatory inhibitory balance: synapses, circuits and systems plasticity*. Kluwer Academic/Plenum, New York, pp 215–228
- Gao B, Fritschy JM, Benke D, Möhler H (1993) Neuron-specific expression of GABA_A receptor subtypes: differential associations of the α_1 - and α_3 -subunits with serotonergic and GABAergic neurons. *Neuroscience* 54:881–892
- Geiger JR, Lubke J, Roth A, Frotscher M, Jonas P (1997) Submillisecond AMPA receptor-mediated signalling at a principal neuron-interneuron synapse. *Neuron* 18:1009–1023
- Griebel G, Perrault G, Simiand J, Cohen C, Granger P, Depoortere H, Francon D, Avenet P, Schoemaker H, Evanno Y, et al (2003) SL651498, a GABA_A receptor agonist with subtype-selective efficacy, as a potential treatment for generalized anxiety disorder and muscle spasms. *CNS Drug Rev* 9:3–20
- Gupta A, Wang Y, Markam H (2000) Organizing principles for a diversity of GABAergic interneurons and synapses in the neocortex. *Science* 287:273–278
- Haefeli W, Martin JR, Schoch P (1990) Novel anxiolytics that act as partial agonists at benzodiazepine receptors. *Trends Pharmacol Sci* 11:452–456
- Harris KD, Henze DA, Hirase H, Leinekugel X, Dragoi G, Czurko A, Buzsaki G (2002) Spike train dynamics predicts theta-related phase precession in hippocampal pyramidal cells. *Nature* 417:738–741
- Hauser J, Rudolph U, Keist R, Möhler H, Feldon J, Yee B (2005) Hippocampal α_5 subunit containing GABA_A receptors modulate expression of prepulse inhibition. *Mol Psychiatry* 10:201–207
- Hensch TK (2005) Critical period plasticity in local cortical circuits. *Nature Rev Neurosci* 6:877–888
- Hensch TK, Stryker MP (2004) Columnar architecture sculpted by GABA circuits in developing cat visual cortex. *Science* 303:1678–1681
- Huckle R (2004) Gaboxadol Lundbeck/Merck. *Curr Opin Investig Drugs* 5:766–773
- Huntsmann MM, Porcello DM, Homanics GE, DeLorey TM, Huguenard JR (1999) Reciprocal inhibitory connections and network synchrony in the mammalian thalamus. *Science* 283:541–543
- Hutcheon B, Morley P, Poulter MO (2000) Developmental change in GABA_A receptor desensitization kinetics and its role in synapse function in rat cortical neurons. *J Physiol (Lond)* 522:3–17
- Jüttner R, Meier J, Grantyn R (2001) Slow IPSC kinetics, low levels of α_1 subunit expression and paired-pulse depression are distinct properties of neonatal inhibitory GABAergic synaptic connections in the mouse superior colliculus. *Eur J Neurosci* 13:2088–2098
- Jurd R, Arras M, Lambert S, Drexler B, Siegwart R, Crestani F, Zaugg M, Vogt KE, Ledermann B, Antkowiak B, et al (2003) General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA(A) receptor β_3 subunit. *FASEB J* 17:250–252
- Kandler K (2004) Activity-dependent organization of inhibitory circuits: lessons from the auditory system. *Curr Opin Neurobiol* 14:96–104
- Kathuria S, Gaetani S, Fegley D, Valino F, Duranti A, Tontini A, Mor M, Tarzia G, La Rana G, Calignano A, Giustino A, Tattoli M, Palmery M, Cuomo V, Piomelli D (2003) Modulation of anxiety through blockade of anandamide hydrolysis. *Nat Med* 9:76–81
- Katona I, Sperlagh B, Sik A, Káfalvi A, Vizi ES, Mackie K, Freund TF (1999) Presynaptically located CB1 cannabinoid receptors regulate GABA release from axon terminals of specific hippocampal interneurons. *J Neurosci* 19:4544–4558
- Katona I, Rancz EA, Acsády L, Ledent C, Mackie K, Hajos N, Freund TF (2001) Distribution of CB1 cannabinoid receptors in the amygdala and their role in the control of GABAergic transmission. *J Neurosci* 21:9506–9518
- Klausberger T, Roberts JD, Somogyi P (2002) Cell type- and input-specific differences in the number and subtypes of synaptic GABA_A receptors in the hippocampus. *J Neurosci* 22:2513–2521
- Klausberger T, Magill PJ, Marton LF, Roberts JDB, Cobden PM, Buzsaki G, Somogyi P (2003) Brain state- and cell type-specific firing of hippocampal interneurons in vivo. *Nature* 421:844–848
- Kopp C, Rudolph U, Tobler I (2004a) Sleep EEG changes after zolpidem in mice. *Neuroreport* 15:2299–2302
- Kopp C, Rudolph U, Löw K, Tobler I (2004b) Modulation of rhythmic brain activity by diazepam: GABA(A) receptor subtype and state specificity. *Proc Natl Acad Sci USA* 101:3674–3679
- Krasowski MD, Koltchine VV, Rick CE, Ye Q, Finn SE, Harrison NL (1998) Propofol and other intravenous anesthetics have sites of action on the gamma-aminobutyric acid type A receptor distinct from that for isoflurane. *Mol Pharmacol* 53:530–538

- Kreitzer AC, Regehr WG (2001a) Retrograde inhibition of presynaptic calcium influx by endogenous cannabinoids at excitatory synapses onto Purkinje cells. *Neuron* 29:717–727
- Kreitzer AC, Regehr WG (2001b) Cerebellar depolarization-induced suppression of inhibition is mediated by endogenous cannabinoids. *J Neurosci* 21:RC174
- Lambert S, Arras M, Vogt KE, Rudolph U (2005) Isoflurane-induced surgical tolerance mediated only in part by beta3-containing GABA(A) receptors. *Eur J Pharmacol* 516:23–27
- Lancel M, Steiger A (1999) Sleep and its modulation by drugs that affect GABA_A receptor function. *Angew Chem Int Ed* 111:2852–2864
- Langen B, Egerland U, Bernoster K, Dost R, Unverferth K, Rundfeldt C (2005) Characterization in rats of the anxiolytic potential of ELB139 [1-(4-chlorophenyl)-4-piperidin-1-yl-1,5-dihydro-imidazol-2-on], a new agonist at the benzodiazepine binding site of the GABA_A receptor. *J Pharmacol Exp Ther* 314:717–724
- Lewis DA, Hashimoto T, Volk DW (2005) Cortical inhibitory neurons and schizophrenia. *Nat Rev Neurosci* 6:312–324
- Liao M, Sonner JM, Jurd R, Rudolph U, Borghese CM, Harris RA, Laster MJ, Eger EI 2nd (2005) Beta3-containing gamma-aminobutyric acid receptors are not major targets for the amnesic and immobilizing actions of isoflurane. *Anesth Analg* 101:412–418
- Lippa A, Czobor P, Stark J, Beer B, Kostakis E, Gravielle M, Bandyopadhyay S, Russek SJ, Gibbs TT, Farb DH, Skolnick P (2005) Selective anxiolysis produced by ocinaplon, a GABA(A) receptor modulator. *Proc Natl Acad Sci USA* 102:7380–7385
- Llano I, Leresche N, Marty A (1991) Calcium entry increases the sensitivity of cerebellar Purkinje cells to applied GABA and decreases inhibitory synaptic currents. *Neuron* 6:565–574
- Löw K, Crestani F, Keist R, Benke D, Brunig I, Benson JA, Fritschy JM, Rulicke T, Bluethmann H, Möhler H, Rudolph U (2000) Molecular and neuronal substrate for the selective attenuation of anxiety. *Science* 290:131–134
- Maejima T, Ohno-Shosaku T, Kano M (2001) Endogenous cannabinoid mediate retrograde signals from depolarized postsynaptic neurons to presynaptic terminals. *Neuron* 29:729–738
- Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2004) Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* 10:793–807
- Marowsky A, Fritschy JM, Vogt KE (2004) Functional mapping of GABA_A receptor subtypes in the amygdala. *Eur J Neurosci* 20:1281–1289
- Marsicano G, Wotjak CT, Azad SC, Bisogno T, Rammes G, Cascio MG, Hermann H, Tang J, Hofmann C, Zieglansberger W, Di Marzo V, Lutz B (2002) The endogenous cannabinoid system controls extinction of aversive memories. *Nature* 418:530–534
- Martina M, Schultz JH, Ehmke H, Monyer H, Jonas P (1998) Functional and molecular differences between voltage-gated K⁺ channels of fast-spiking interneurons and pyramidal neurons of rat hippocampus. *J Neurosci* 18:1811–1825
- McKernan RM, Rosahl TW, Reynolds DS, Sur C, Wafford KA, Atack JR, Farrar S, Myers J, Cook G, Ferris P, Garrett L, Bristow L, Marshall G, Macaulay A, Brown N, Howell O, Moore KW, Carling RW, Street LJ, Castro JL, Ragan CI, Dawson GR, Whiting PJ (2000) Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA_A receptor α_1 subtype. *Nat Neurosci* 3:587–592
- Metha MR, Lee AK, Wilson MA (2002) Role of experience and oscillations in transforming a rate code into a temporal code. *Nature* 417:741–746
- Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, et al (1997) Sites of alcohol and volatile anaesthetic action on GABA (A) and glycine receptors. *Nature* 389:385–389
- Mody I, Pearce RA (2004) Diversity of inhibitory neurotransmission through GABA(A) receptors. *Trends Neurosci* 27:569–575
- Möhler H (2001) Functions of GABA receptors: pharmacology and pathophysiology. In: Möhler H (ed) *Pharmacology of GABA and glycine neurotransmission*. Springer, Berlin Heidelberg New York, pp 101–116
- Möhler H (2002) Pathophysiological aspects of diversity in neuronal inhibition: a new benzodiazepine pharmacology. *Dialogues Clin Neurosci* 4:261–269
- Möhler H, Benke D, Fritschy JM, Benson J (2000) The benzodiazepine site of GABA_A receptors. In: Martin DL, Olsen RW (eds) *GABA in the nervous system: the view at fifty years*. Lippincott, Philadelphia, pp 97–112
- Möhler H, Fritschy JM, Rudolph U (2002) A new benzodiazepine pharmacology. *J Pharm Exptl Ther* 300:2–8
- Möhler H, Fritschy JM, Vogt K, Crestani F, Rudolph U (2005) Pathophysiology and pharmacology of GABA_A receptors. In: Holsboer F, Ströhle A (eds) *Anxiety and anxiolytic drugs. Handbook of experimental pharmacology*, vol 169. Springer, Berlin Heidelberg New York, pp 225–247
- Monyer H, Markram H (2004) Interneuron Diversity series: Molecular and genetic tools to study GABAergic interneuron diversity and function. *Trends Neurosci* 27:90–97
- Moss SJ, Smart TG (2001) Constructing inhibitory synapses. *Nat Rev Neurosci* 2:240–250
- Navarro JF, Buron E, Martin-Lopez M (2002) Anxiogenic-like activity of L-655,708, a selective ligand for the benzodiazepine site of GABA(A) receptors which contain the alpha-5 subunit, in the elevated plus-maze test. *Prog Neuropsychopharmacol Biol Psychiatry* 26:1389–1392
- Navarro JF, Buron E and Martin-Lopez M (2004) Behavioral profile of L-655 708, a selective ligand for the benzodiazepine site of GABA_A receptors which contain the α_5 subunit in social encounters between male mice. *Aggress Behav* 30:319–325
- Nusser Z, Sieghart W, Stephenson FA, Somogyi P (1996) The α_6 subunit of the GABA_A receptor is concentrated in both inhibitory and excitatory synapses on cerebellar granule cells. *J Neurosci* 16:103–114
- Nusser Z, Sieghart W and Somogyi P (1998) Segregation of different GABAA receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci* 18:1693–1703
- Nyíri G, Freund TF and Somogyi P (2001) Input-dependent synaptic targeting of α_2 subunit containing GABA_A receptors in hippocampal pyramidal cells of the rat. *Eur J Neurosci* 13:428–442
- O'Keefe J, Nadel L (1978) *The hippocampus as a cognitive map*. Clarendon, Oxford, pp 477–543
- O'Keefe J, Recce ML (1993) Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus* 3:317–330
- Paulsen O, Moser EI (1998) A model of hippocampal memory encoding and retrieval: GABAergic control of synaptic plasticity. *Trends Neurosci* 21:273–278
- Pawelzik H, Hughes DI, Thomson AM (2002) Physiological and morphological diversity of immunocytochemically defined parvalbumin- and cholecystokinin-positive interneurons in CA1 of the adult rat hippocampus. *J Comp Neurol* 443:346–367
- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G (2000) GABA_A receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 101:815–850
- Pitler TA, Alger BE (1992) Postsynaptic spike firing reduces synaptic GABA_A responses in hippocampal pyramidal cells. *J Neurosci* 12:4122–4132
- Pitler TA, Alger BE (1994) Depolarization-induced suppression of GABAergic inhibition in rat hippocampal pyramidal cells: G protein involvement in a presynaptic mechanism. *Neuron* 13:1447–1455

- Pörtl A, Hauer B, Fuchs K, Tretter V, Sieghart W (2003) Subunit composition and quantitative importance of GABA_A receptors subtypes in the cerebellum of mouse and rat. *J Neurochem* 87:1444–1455
- Represa A, Ben-Ari Y (2005) Trophic actions of GABA on neuronal development. *Trends Neurosci* 28:278–283
- Reynolds DS, Rosahl TW, Cirone J, O'Meara GF, Haythornthwaite A, Newman RJ, Myers J, Sur C, Howell O, Rutter AR, et al (2003) Sedation and anesthesia mediated by distinct GABA(A) receptor isoforms. *J Neurosci* 23:8608–8617
- Rijnsoever C van, Tauber M, Choulli MK, Keist R, Rudolph U, Möhler H, Fritschy JM, Crestani F (2004) Requirement of α_5 GABA_A receptors for the development of tolerance to the sedative action of diazepam in mice. *J Neurosci* 24:6785–6790
- Rudolph U, Antkowiak B (2004) Molecular and neuronal substrates for general anaesthetics. *Nat Rev Neurosci* 5:709–720
- Rudolph U, Möhler H (2004) Analysis of GABA_A receptor function and dissection of pharmacology of benzodiazepines and general anaesthetics by mouse genetics. *Annu Rev Pharmacol Toxicol* 44:475–498
- Rudolph U, Möhler H (2006) GABA-based therapeutic approaches: GABA_A receptor subtype functions. *Curr Opin Pharmacol* 6:18–23
- Rudolph U, Crestani F, Benke D, Brünig I, Benson J, Fritschy JM, Martin JR, Bluethmann H, Möhler H (1999) Benzodiazepine actions mediated by specific γ -aminobutyric acid_A receptor subtypes. *Nature* 401:796–800
- Sieghart W, Sperk G (2002) Subunit composition, distribution and function of GABA_A receptor subtypes. *Curr Top Med Chem* 2:795–816
- Skaggs WE, McNaughton BL, Wilson MA, Barnes CA (1996) Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* 6:149–172
- Sternfeld F, Carling RW, Jelley RA, Ladduwahetty T, Merchant KJ, Moore KW, Reeve AJ, Street LJ, O'Connor D, Sohal B, et al (2004) Selective, orally active gamma-aminobutyric acid_A α_5 receptor inverse agonists as cognition enhancers. *J Med Chem* 47:2176–2179
- Storustovu S, Ebert B (2003) Gaboxadol: in vitro interaction studies with benzodiazepines and ethanol suggest functional selectivity. *Eur J Pharmacol* 467:49–56
- Tobler I, Kopp C, Deboer T, Rudolph U (2001) Diazepam-induced changes in sleep: role of the α_1 GABA_A receptor subtype. *Proc Natl Acad Sci USA* 98:6464–6469
- Traub RD, Draguhn A, Whittington MA, Baldeweg T, Bibbig A, Buhl EH, Schmitz D (2002) Axonal gap junctions between principal neurons: a novel source of network oscillations, and perhaps epileptogenesis. *Rev Neurosci* 13:1–30
- Vicini S, Ferguson C, Prybylowski K, Kralic J, Morrow AL, Homanics GE (2001) GABA_A receptor α_1 subunit deletion prevents developmental changes of inhibitory synaptic currents in cerebellar neurons. *J Neurosci* 21:3009–3016
- Vincent P, Marty A (1993) Neighboring cerebellar Purkinje cell communicate via retrograde inhibition of common presynaptic interneurons. *Neuron* 11:885–893
- Wallner M, Hancher HJ, Olsen RW (2003) Ethanol enhances α_4 β_3 δ and α_6 β_3 δ gamma-aminobutyric acid type A receptors at low concentration known to affect humans. *Proc Natl Acad Sci USA* 100:15218–15223
- Whiting PJ (2003) The GABA_A receptor gene family: new opportunities for drug development. *Curr Opin Drug Discov Dev* 6:648–655
- Whiting P, Wafford KA, McKernan RM (2000) Pharmacologic subtypes of GABA_A receptors based on subunit composition In: Martin DL, Olsen RW (eds) *GABA in the nervous system: the view at fifty years*. Lippincott, Philadelphia, pp 113–126
- Wiesel TN, Hubel DH (1963) Single cell responses in striate cortex of kittens deprived of vision in one eye. *J Neurophysiol* 26:1003–1017
- Wilson RI, Nicoll RA (2001) Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. *Nature* 410:588–592 [erratum appears in *Nature* 2001 411:974]
- Yee BK, Hauser J, Dolgov VV, Keist R, Möhler H, Rudolph U, Feldon J (2004) GABA receptors containing the α_5 subunit mediate the trance effect in aversive and appetitive conditioning and extinction of conditioned fear. *Eur J Neurosci* 20:1928–1936
- Yee BK, et al (2005) A schizophrenia-related sensorimotor deficit links α_3 -containing GABA_A receptors to a dopamine hyperfunction. *Proc Natl Acad Sci USA* 102:17154–17159
- Zeller A, Arras M, Lazaris A, Jurd R, Rudolph U (2005) Distinct molecular targets for the central respiratory and cardiac actions of the general anesthetics etomidate and propofol. *FASEB J* 12:1677–1679